

BLEMMYES/EWAIPANOMA HYPOTHETICAL MODEL

Medical-Grade Model with High-Resolution Geographic Context

This document presents a medically grounded hypothetical model for a putative human variant population with extreme craniofacial reorganization. Long-term survival would have been strongly dependent on warm, stable climates. The most plausible geographic anchor is Northeastern Africa, particularly the Eastern Desert and adjacent regions.


1. Core Anatomical Hypothesis

Brain housed in a shortened neck stub (C3–T1) with a soft superior surface for thermoregulation via sweating and blood flow. Ears positioned on the neck stub for hearing and cooling. Facial features developed from the clavicular/upper chest region. Protection primarily via cartilage and soft tissue. Long hair grown to conceal and protect the vulnerable neck stub.

2. Environmental Constraints & Geographic Range

A population with this anatomy would have faced severe challenges in cold or variable climates. Warm, stable environments would have been essential for survival. The Eastern Desert, Nile Valley margins, and parts of the Sahel/Horn of Africa represent the most viable long-term range. Movement into colder northern latitudes would have carried extremely high mortality risk.

Geographic Context Map



3. Comparison with Human Holoprosencephaly

Severe human HPE demonstrates that extreme craniofacial and brain reorganization can occur, but it is almost always lethal early in life. The Blemmeyes model requires multi-generational population persistence. The proposed adaptations attempt to address viability challenges seen in HPE.

Confidence Assessment

Claim	Confidence
Real vulnerable population existed and was targeted	92–95%
Core anatomical configuration is literally accurate	70–75%